

## VACCINES REQUIRED FOR CHILD CARE/PRESCHOOL ATTENDANCE

July 1, 2016 – June 30, 2017

	<b>Hepatitis B</b>	<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)	<b>Hib</b> ( <i>Haemophilus influenzae</i> type B)	<b>Polio</b>	<b>PCV</b> (Pneumococcal Conjugate)	<b>MMR</b> (Measles, Mumps, Rubella)	<b>Varicella</b> (Chickenpox)
<b>By 3 Months</b> (on or before last day of mo 2)	<b>2 doses</b> May get Dose 1 at birth and Dose 2 as early as 1 month of age	<b>1 dose</b>	<b>1 dose</b>	<b>1 dose</b>	<b>1 dose</b>	<b>Not given before 12 months of age</b>	<b>Not given before 12 months of age</b>
<b>By 5 Months</b> (on or before last day of mo 4)	<b>2 doses</b>	<b>2 doses</b>	<b>2 doses</b>	<b>2 doses</b> May get Dose 2 as early as 4 months of age	<b>2 doses</b>		
<b>By 7 Months</b> (on or before last day of mo 6)	<b>2 doses</b>	<b>3 doses</b> May get Dose 3 as early as 6 months of age	<b>3 doses</b>	<b>2 doses</b>	<b>3 doses</b>		
<b>By 16 Months</b> (on or before last day of mo 15)	<b>2 doses</b>	<b>3 doses</b>	<b>4 doses</b>	<b>2 doses</b>	<b>4 doses*</b>	<b>1 dose</b> May get Dose 1 as early as 12 months of age	<b>1 dose</b> May get Dose 1 as early as 12 months of age <b>OR</b> <b>Healthcare provider verified disease</b>
<b>By 19 Months</b> (on or before last day of mo 18)	<b>3 doses</b>	<b>4 doses</b> May get Dose 4 as early as 12 months as long as 6 months separate Dose 3 and Dose 4	<b>4 doses</b>	<b>3 doses</b>	<b>4 doses*</b>	<b>1 dose</b>	<b>1 dose</b> <b>OR</b> <b>Healthcare provider verified disease</b>
<b>By 7 Years</b> (on or before last day of year 6) or by <b>Kindergarten Entry</b>	<b>3 doses</b>	<b>5 doses</b>	<b>Not given after 5 years of age unless child has medical condition</b>	<b>4 doses</b>	<b>Not given after 5 years of age unless child has medical condition</b>	<b>2 doses</b>	<b>2 doses</b> <b>OR</b> <b>Healthcare provider verified disease</b>

\*Some children may get 5 total doses. A single supplemental dose of PCV13 is recommended, but not required, for all children aged 14–59 months who got 4 doses of PCV7.

- School-aged children (K-12) in before and after-school programs must meet the immunization requirements for their grade in school.
- Find information on other vaccines recommended, but not required, for child care/preschool attendance: [www.immunize.org/cdc/schedules/](http://www.immunize.org/cdc/schedules/)
- Review the Individual Vaccine Requirements Summary for more detailed information: [www.doh.wa.gov/Immunization/schoolandchildcare/VaccineRequirements.aspx](http://www.doh.wa.gov/Immunization/schoolandchildcare/VaccineRequirements.aspx)

## Minimum Age & Interval for Valid Vaccine Doses

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
Hepatitis B (HepB)	Dose 1	Birth	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>The final dose in the series should be given at least 24 weeks of age.</li> </ul>
	Dose 2	4 weeks	8 weeks between Dose 2 & 3	
	Dose 3	24 weeks	16 weeks between Dose 1 & 3	
Diphtheria, Tetanus, and Pertussis (DTaP/DT)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Typical vaccine schedule: 2, 4, 6, and 15-18 months of age.</li> <li>Recommended: 6 months between Dose 3 and Dose 4, but at least 4 months minimum interval acceptable.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	12 months	6 months between Dose 4 & 5	
	Dose 5	4 years	--	
<i>Haemophilus influenzae</i> type B (Hib)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>If all 3 doses of PedvaxHIB given, only need 3 doses total.</li> <li>Only one dose required if the dose given on or after 15 months of age. Review the Individual Vaccine Requirements Summary for minimum doses required: <a href="http://www.doh.wa.gov/immunization/schoolandchildcare/VaccineRequirements.aspx">www.doh.wa.gov/immunization/schoolandchildcare/VaccineRequirements.aspx</a></li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	8 weeks between Dose 3 & 4	
	Dose 4	12 months	--	
Pneumococcal Conjugate (PCV7 or PCV13)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>A single supplemental dose of PCV13 recommended for all children 14–59 months of age who got 4 doses of PCV7.</li> <li>Only one dose required if the dose given on or after 24 months of age. Review the Individual Vaccine Requirements Summary for minimum doses required: <a href="http://www.doh.wa.gov/immunization/schoolandchildcare/VaccineRequirements.aspx">www.doh.wa.gov/immunization/schoolandchildcare/VaccineRequirements.aspx</a></li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	8 weeks between Dose 3 & 4	
	Dose 4	12 months	--	
Polio (IPV or OPV)	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Three doses acceptable if child got Dose 3 on or after the 4<sup>th</sup> birthday.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	4 years	--	
Measles, Mumps, and Rubella (MMR or MMRV)	Dose 1	12 months	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>MMRV (MMR + varicella) may be used in place of separate MMR and varicella vaccines.</li> <li>Must get the same day as VAR <b>OR</b> at least 28 days apart.</li> <li>4-day grace <b>DOES</b> apply between doses of the same live vaccine such as MMR/MMR or MMRV/MMRV. The 4 day grace period <b>DOES NOT</b> apply between Dose 1 and Dose 2 of different live vaccines, such as between MMR and Varicella or between MMR and live flu vaccine.</li> </ul>
	Dose 2	13 months	--	
Varicella (chickenpox) (VAR)	Dose 1	12 months	3 months between Dose 1 & 2 (12 months through 12 years) 4 weeks between Dose 1 & 2 (13 years and older)	<ul style="list-style-type: none"> <li>Recommended: 3 months between varicella doses, but at least 28 days minimum interval acceptable. Minimum age of 13 months acceptable.</li> <li>Must get the same day as MMR <b>OR</b> at least 28 days apart.</li> <li>4-day grace <b>DOES</b> apply between doses of the same live vaccine; <b>DOES NOT</b> apply between doses of different live vaccines.</li> </ul>
	Dose 2	15 months	--	